
**GARFIELD COUNTY
2019 AIR QUALITY MONITORING REPORT**

Prepared for:

**Garfield County Public Health Department
195 West 14th Street
Rifle, CO 81650
and
2014 Blake Ave.
Glenwood Springs, CO 81601**

Prepared by



1901 Sharp Point Dr., Suite F
Fort Collins, CO 80525
970-484-7941
www.air-resource.com

June 30, 2020

A decorative graphic in the bottom left corner consisting of three stylized mountain peaks in shades of green and blue.

TABLE OF CONTENTS

<u>Section</u>	<u>Page</u>
LIST OF ACRONYMS AND ABBREVIATIONS	iv
EXECUTIVE SUMMARY	v
1.0 INTRODUCTION	1-1
2.0 METEOROLOGICAL SUMMARIES	2-1
3.0 CRITERIA POLLUTANT SUMMARIES	3-1
3.1 Ozone	3-2
3.2 Nitrogen Dioxide	3-6
3.3 Particulate Matter	3-7
3.3.1 PM ₁₀ Measurements	3-8
3.3.2 PM _{2.5} Measurements	3-9
4.0 VOLATILE ORGANIC COMPOUNDS	4-1
4.1 Continuous Methane and Non-Methane Hydrocarbons	4-1
4.2 Speciated Non-Methane Organic Compounds	4-3
4.3 Carbonyls	4-7
4.4 Hazardous Air Pollutants (HAPS)	4-9
4.4.1 Annual Average HAP Concentrations	4-9
5.0 REFERENCES	5-1
APPENDIX A Quarterly Time Series Plots	A-1
APPENDIX B SNMOC Concentrations	B-1
APPENDIX C Carbonyl Concentrations	C-1

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
Figure 1-1. Map of 2019 Garfield County air quality monitoring sites.	1-2
Figure 2-1. Map with wind roses depicting 2019 wind speed and direction measured at the Garfield County air quality monitoring sites.	2-1
Figure 3-1. Fourth highest 8-Hour ozone averages measured at all Garfield County sites in comparison to the NAAQS.	3-3
Figure 3-2. Fourth highest 8-Hour ozone averages measured at the Rifle site in comparison to the NAAQS. Note that an exceedance of the standard is not considered a violation unless the annual fourth-highest daily maximum 8-hr concentration averaged over three years exceeds 70 ppb.	3-3
Figure 3-3. Fourth highest 8-Hour ozone averages measured at the mobile monitor in comparison to the NAAQS.	3-4
Figure 3-4. Fourth highest 8-Hour ozone averages measured at the Carbondale site in comparison to the NAAQS.	3-4
Figure 3-5. Daily maximum 8-hour averages of ozone monitored at the Rifle site in 2019 in comparison to the NAAQS.	3-4
Figure 3-6. Daily maximum 8-hour averages of ozone monitored at the mobile monitor (Battlement Mesa site) in 2019 in comparison to the NAAQS.	3-4
Figure 3-7. Daily maximum 8-hour averages of ozone monitored at the Carbondale site in 2019 in comparison to the NAAQS.	3-5
Figure 3-8. Annual average nitrogen dioxide monitored at the mobile monitor (Battlement Mesa/Glenwood Springs) in 2019 in comparison to the NAAQS.	3-6
Figure 3-9. Daily maximum 1-hour averages of nitrogen dioxide monitored at the mobile monitor (Battlement Mesa site) in 2019 in comparison to the NAAQS.	3-7
Figure 3-10. Annual average PM ₁₀ measured at the Mobile site.	3-8
Figure 3-11. Annual average PM _{2.5} measured at the Garfield County sites in comparison to the NAAQS.	3-9
Figure 3-12. Highest and 98th percentile 24-hour average PM _{2.5} measured at the Carbondale site.	3-10
Figure 3-13. Highest and 98th percentile 24-hour average PM _{2.5} measured at the Rifle site. Note that exceedances of the NAAQS are not considered a violation until the 3-year average of the 98 th percentile value exceeds the NAAQS.	3-10
Figure 3-14. Highest and 98th percentile 24-hour average PM _{2.5} measured at the mobile monitor (Battlement Mesa/Glenwood Springs) in comparison to the NAAQS.	3-11

LIST OF FIGURES (continued)

<u>Figure</u>		<u>Page</u>
Figure 4-1.	Daily average methane measured at the mobile monitor (Battlement Mesa site) in 2019.	4-2
Figure 4-2.	Daily average canister samples and continuous measurements of non-methane hydrocarbons measured at the mobile monitor (Battlement Mesa site) in 2019.	4-2
Figure 4-3	Average SNMOC concentrations measured at Garfield County sites between 2008 and 2019.	4-4
Figure 4-4.	2019 24-hour SNMOC measurements by category in units of ppbV.	4-5
Figure 4-5.	2019 24-hour SNMOC measurements by category in units of ppbC.	4-6
Figure 4-6.	Average carbonyl concentrations measured at Garfield County sites between 2008 and 2019.	4-7
Figure 4-7.	2019 24-hour major carbonyl compound concentrations in units of ppbV.	4-8

LIST OF TABLES

<u>Table</u>		<u>Page</u>
Table 1-1	Garfield County Parameters Monitored by Site	1-3
Table 4-1	Parachute Site Annual Average Mass Trends (HAPs Parameters) 2008-2019	4-11
Table 4-2	Rifle Site Annual Average Mass Trends (HAPs Parameters) 2008-2019	4-12
Table 4-3	Bell Ranch Site Annual Average Mass Trends (HAPs Parameters) 2008-2019	4-13
Table 4-4	Mobile Monitor – Battlement Mesa Site Annual Average Mass Trends (HAPs Parameters) 2008-2019	4-14
Table 4-5	Carbondale Site Annual Average Mass Trends (HAPs Parameters) 2008-2019	4-15

LIST OF ACRONYMS AND ABBREVIATIONS

ARS	Air Resource Specialists, Inc.
ATSDR	Agency for Toxic Substances and Disease Registry
BMCO	Battlement Mesa, Colorado Air Quality Monitoring Site
BRCO	Bell Ranch, Colorado Air Quality Monitoring Site
BTEX	Benzene, toluene, ethylbenzene, and xylenes
CDPHE	Colorado Department of Public Health and Environment
CH ₄	Methane
EPA	U.S. Environmental Protection Agency
ERG	Eastern Research Group, Inc.
FRM	Federal Reference Method
GCPH	Garfield County Public Health
GHG	Greenhouse Gas
GSCO	Glenwood Springs, Colorado Air Quality Monitoring Site
HAPs	Hazardous Air Pollutants
NAAQS	National Ambient Air Quality Standards
NATTS	National Air Toxics Trends Stations
NIOSH	National Institute for Occupational Safety and Health
NMOC	Non-Methane Organic Compounds
NO	Nitric Oxide
NO ₂	Nitrogen Dioxide
NO _x	Oxides of nitrogen
O ₃	Ozone
PACO	Parachute, Colorado Air Quality Monitoring Site
PAH	Polycyclic aromatic hydrocarbons
PM _{2.5}	Particulate matter \leq 2.5 microns in diameter
PM ₁₀	Particulate matter \leq 10 microns in diameter
PPB	Parts per billion
PPM	Parts per million
PPBC	Parts per billion carbon
RECO	Rifle, Colorado Air Quality Monitoring Site (formerly RICO)
RFCO	Carbondale, Colorado (Roaring Fork) Air Quality Monitoring Site
SNMOC	Speciated non-methane organic compounds
THC	Total Hydrocarbons
TNMOC	Total non-methane organic compounds
UATMP	Urban Air Toxics Monitoring Program
VOC	Volatile organic compounds

EXECUTIVE SUMMARY

This report summarizes air quality monitoring data collected through 2019 in Garfield County, Colorado. Ambient air quality was monitored at five locations in the County in 2019. Collected pollutants and parameters are described below.

Criteria Air Pollutants are subject to National Ambient Air Quality Standards (NAAQS) under the Clean Air Act. Air quality standards are based on human health and/or environmental impacts.

Parameter	NAAQS	Parameter Description
Ozone (O₃)	Rolling 8-hour average not to exceed 70 ppb	Ozone is not emitted directly into the air, but is formed by chemical reactions between oxides of nitrogen (NO _x) and volatile organic compounds (VOC) in the presence of sunlight. Emissions from industrial facilities and electric utilities, motor vehicle exhaust, gasoline vapors, and chemical solvents are some of the major sources of NO _x and VOCs. Breathing ozone can trigger a variety of health problems, particularly for children, the elderly, and people of all ages who have lung diseases such as asthma. Ground level ozone can also have harmful effects on sensitive vegetation and ecosystems
Nitrogen Dioxide (NO₂)	Annual average not to exceed 53 ppb or 1-hour average not to exceed 100 ppb	Nitrogen dioxide (NO ₂) is one of a group of highly reactive gasses known as "oxides of nitrogen," or "nitrogen oxides (NO _x)." NO ₂ forms quickly from emissions from cars, trucks, and buses, power plants, and off-road equipment. In addition to contributing to the formation of ground-level ozone and fine particle pollution, NO ₂ is linked with a number of adverse effects on the respiratory system.
Particulate Matter (PM₁₀) and (PM_{2.5})	24-hour PM ₁₀ average not to exceed 150 µg/m ³ Annual PM _{2.5} average not to exceed 12 µg/m ³ and/or 24-hour average not to exceed 35 µg/m ³	Particulate Matter describes a mixture of solid particles and liquid droplets found in the air. Some particles, such as dust, dirt, soot, or smoke, are large or dark enough to be seen with the naked eye. Others are so small they can only be detected using an electron microscope. Types of particulate matter include: <ul style="list-style-type: none"> • PM₁₀: inhalable particles, with diameters that are generally 10 micrometers and smaller; and • PM_{2.5}: fine inhalable particles, with diameters that are generally 2.5 micrometers and smaller.

Air Toxics are gaseous, aerosol or particle pollutants present in the air in varying concentrations with characteristics such as toxicity or persistence that can be hazardous to human, plant or animal life. These compounds include Volatile Organic Compounds, (VOCs), including methane (CH₄), subsets of speciated non-methane organic compounds (SNMOCs), carbonyls, and hazardous air pollutants (HAPs).

Parameter	NAAQS	Parameter Description
VOC	No ambient air quality standards exist for toxic air pollutants	Volatile organic compounds (VOC) means any compound of carbon, excluding carbon monoxide, carbon dioxide, carbonic acid, metallic carbides or carbonates, and ammonium carbonate, which participates in atmospheric photochemical reactions.
SNMOC		Identical to VOCs with methane excluded, including compounds associated with natural gas and gasoline vapors, crude oil products, and gas-fired engines.
CARBONYLS		Carbonyl compounds include formaldehyde, aldehydes and ketones. Formaldehyde is a major compound in the formation of photochemical ozone. Short term exposure is known to cause irritation of the eyes, skin, and mucous membranes of the respiratory tract.
HAP		Hazardous air pollutants are those pollutants that cause or may cause cancer or other serious health effects, such as reproductive effects or birth defects, or adverse environmental and ecological effects

Meteorological data consist of physical parameters that are measured directly by instrumentation. The following data are collected to better understand the local conditions and transport of air pollutants:

- Wind speed
- Wind direction
- Ambient Temperature
- Relative Humidity
- Barometric Pressure
- Precipitation

Monitoring results through 2019 indicate that:

- Air quality measurements in Garfield County did not violate NAAQS for PM₁₀, PM_{2.5}, O₃, or NO₂.
- Light alkanes (e.g., ethane, propane, iso/n-butane, and iso/n-pentane), which are commonly associated with natural gas, made up between 68% and 88% of the total SNMOC compounds measured at all air monitoring sites. These light alkanes may contribute to ozone formation and odor issues but are not considered HAPs.
- SNMOC compounds concentrations increased slightly at the Rifle, Bell Ranch, and Carbondale sites. Carbonyl concentrations decreased at all sites.

Detailed air quality monitoring summaries for data collected through 2019 are presented in this report. Additional information, including real-time air quality data, previous air quality data reports, educational materials, air quality management plans, emissions inventories, and health risk assessments are available on the Garfield County Air Quality Management website (<http://www.garfield-county.com/air-quality/>). Any questions regarding this report or air quality in Garfield County should be addressed to:

Garfield County Public Health
195 West 14th Street
Rifle, CO 81650

And

2014 Blake Ave
Glenwood Springs, CO 81601

Phone: 970-625-5200

1.0 INTRODUCTION

Garfield County Public Health (GCPH) has implemented and maintains a network of air quality monitors designed to serve a wide range of purposes related to public health and environmental concerns. This annual air quality data report presents air quality data collected in Garfield County through 2019. The 2019 monitoring network in Garfield County consisted of five monitoring locations, as shown on a map in Figure 1-1 and described below.

- Parachute (PACO): Parachute is a small urban center of approximately 1,300 people within very close proximity to natural gas development activities. The town is located along Interstate 70 and is the transportation hub for heavily traveled roads which service natural gas development in the surrounding area.
- Rifle (RECO): Rifle is a growing urban center on the Interstate 70 corridor with an estimated population of approximately 9,500 people. Rifle is in close proximity to oil and gas development activities, and is also central to industrial support for the oil and gas industry. In 2017, the Rifle equipment moved to the roof of the Rifle Library, necessitating a change in the site ID.
- Bell Ranch (BRCO): The Bell Ranch site is a rural homestead approximately four miles south of the town of Silt, in close proximity to natural gas development.
- Battlement Mesa (BMCO): Battlement Mesa is an unincorporated community located about 1.5 miles southeast of Parachute. A VOC sampler was located here in September 2010, and the Garfield County mobile monitoring station was located here in November 2012. This was in response to proposed natural gas development within the community, and to begin developing baseline data in advance of potential large scale production. This mobile monitor was relocated to Glenwood Springs in February 2015, and back to Battlement Mesa in March 2016. Annual average data for the Glenwood Springs site were presented in the 2015 Annual Data report.
- Carbondale (RFCO): Carbondale is a small town located about 12.5 miles southeast of Glenwood Springs at the confluence of the Roaring Fork and Crystal River valleys. The Carbondale, or Roaring Fork, site began monitoring in March 2012.

Table 1-1 lists the pollutants and other parameters monitored at each site. Monitoring at these stations is performed by GCPH with technical support from several agencies, as noted in Table 1-1. Real-time data are displayed on the Garfield County Air Quality website (<http://www.garfield-county.com/air-quality/>).




Figure 1-1. Map of 2019 Garfield County air quality monitoring sites.

Table 1-1
Garfield County
Parameters Monitored by Site

Component	Method	Sampling Frequency	Technical Analysis
Rifle, Colorado			
SNMOC	TO-12	24-hour (1/6 day)	ERG
Carbonyls	TO-11A	24-hour (1/12 day)	ERG
PM _{2.5}	BAM	Hourly	ARS
Ozone	42C	Hourly	ARS
Meteorology	Various	Hourly	ARS
Visibility Web Camera	Digital	15-min	ARS
Parachute, Colorado			
SNMOC	TO-12	24-hour (1/6 day)	ERG
Carbonyls	TO-11A	24-hour (1/12 day)	ERG
Meteorology	Various	Hourly	ARS
Bell Ranch, Colorado			
SNMOC	TO-12	24-hour (1/6 day)	ERG
Carbonyls	TO-11A	24-hour (1/12 day)	ERG
Meteorology	Various	Hourly	ARS
Mobile Monitor (Battlement Mesa, Colorado)*			
SNMOC	TO-12	24-hour (1/6 day)	ERG
Carbonyls	TO-11A	24-hour (1/12 day)	ERG
PM ₁₀	BAM	Hourly	ARS
PM _{2.5}	BAM	Hourly	ARS
Ozone	API	Hourly	ARS
NO/NO ₂ /NO _x	API	Hourly	ARS
CH ₄ /NMHC/THC	Baseline MOCON	Hourly	ARS
Meteorology	Various	Hourly	ARS
Carbondale, Colorado			
SNMOC	TO-12	24-hour (1/12 day)	ERG
PM _{2.5}	E-BAM	Hourly	ARS
Ozone	2B	Hourly	ARS
Meteorology	Various	Hourly	ARS

*The mobile monitor operated in Glenwood Springs February 2015 through February 2016 and returned to Battlement Mesa in March 2016.

2.0 METEOROLOGICAL SUMMARIES

- Meteorological data are collected along with other air quality parameters to better understand the local conditions and transport of air pollutants. Meteorological data collected in Garfield County includes wind speed, wind direction, temperature, relative humidity, barometric pressure and precipitation.

Figure 2-1 presents a map overlaid with wind roses from each of the Garfield County monitoring sites. Wind roses depict wind direction and wind speed. The direction of the bar signifies the direction the wind is coming from, the length of the bars indicates the cumulative frequency from each direction, and the colors indicate wind speed. The map shows that winds at the Garfield County sites are influenced by flow along the Colorado River Basin, where Interstate 70 crosses through the county. Airflow is also influenced by drainage flows through valleys along various Colorado River tributaries.




Figure 2-1. Map with wind roses depicting 2019 wind speed and direction measured at the Garfield County air quality monitoring sites.

3.0 CRITERIA POLLUTANT SUMMARIES

The Clean Air Act requires the Environmental Protection Agency (EPA) to set two types of National Ambient Air Quality Standards (NAAQS) for pollutants that include ground-level ozone (O₃), particle matter (PM_{2.5} and PM₁₀), lead (Pb), nitrogen dioxide (NO₂), carbon monoxide (CO), and sulfur dioxide (SO₂). The types of standards are as follows:



- **Primary Standards:** These standards are designed to protect public health with an adequate margin of safety, including the health of sensitive populations such as asthmatics, children, and the elderly.
- **Secondary Standards:** These standards are designed to protect public welfare from adverse pollution effects, including visibility impairment and effects on the environment (e.g., vegetation, soils, water, and wildlife).

Each standard is defined by an averaging time, a concentration level, and a period of evaluation. Note that a measured value which exceeds the concentration level may be termed an “exceedance”, but the concentration measured is not considered a “violation” of the standard unless all criteria are considered, including the period of evaluation.

Criteria pollutant monitoring in Garfield County includes PM₁₀, PM_{2.5}, O₃ and NO₂. This section includes summaries for each of these pollutants and comparisons to the relevant NAAQS. Note that, to date, air quality measurements in Garfield County have not violated NAAQS for any of the criteria pollutants measured.

3.1 OZONE

Ozone is a secondary pollutant, meaning it is not emitted directly from sources, but is formed from photochemical interactions of volatile organic compounds (VOCs) and oxides of nitrogen (NO_x) in the presence of sunlight. The basic formation and depletion equations for O_3 are presented below:



Without the presence of VOCs, the diurnal cycle is a balanced reaction, with equal production and depletion of O_3 . When VOCs are present, they can react with nitric oxide (NO) to produce NO_2 as follows:



This effectively creates competition for NO, allowing O_3 to build up instead of being depleted by NO. Also, when NO reacts with hydrocarbons, additional NO_2 is produced without consuming O_3 , and the produced NO_2 can further react to produce more O_3 . A diurnal cycle will generally show lowest O_3 concentrations in the early morning hours and maximum concentrations in the late afternoon. This pattern results from daytime photochemical production from NO_x ($\text{NO} + \text{NO}_2$) and VOC precursors, and O_3 loss by dry deposition and reaction with NO at night.

The NAAQS for O_3 was lowered in 2015 by the Environmental Protection Agency (EPA) to 70 ppb (previously 75 ppb) over an 8-hour period. An exceedance of the standard occurs when an 8-hour average O_3 concentration is greater than 70 ppb. A violation of the standard occurs when the three-year average of the fourth highest daily maximum 8-hour average ozone concentration exceeds 70 ppb.

Ozone measurements began in June 2008 at the Rifle site, in March 2012 at the Carbondale site, and in November 2012 at the mobile site in Battlement Mesa (located in Glenwood Springs in 2015). Figures 3-1 through 3-4 show annual trends for O_3 monitored at all Garfield County sites through 2019. At the Rifle site, the highest recorded 8-hour averages exceeded the standard in 2008, 2012, and 2018, but the 3-year average of the 4th highest values did not, so this was not considered a violation of the standard.

The pump in the ozone analyzer at the Carbondale site failed in August 2019. The instrument was returned to ARS for evaluation and repair. Following reinstallation in September the instrument continued to operate erratically. During the third quarter maintenance visit the power supply was replaced and normal data operations resumed.

Figures 3-5 through 3-7 present daily maximum 8-hour averages of O_3 monitored at the Rifle, Mobile, and Carbondale sites, respectively, in 2019.




Figure 3-1. Fourth Highest 8-Hour ozone averages measured at all Garfield County sites in comparison to the NAAQS.




Figure 3-2. Fourth highest and maximum 8-Hour ozone averages measured at the Rifle site in comparison to the NAAQS. Note that an exceedance of the standard is not considered a violation unless the annual fourth-highest daily maximum 8-hr concentration averaged over three years exceeds 70 ppb.




Figure 3-3. Fourth highest and maximum 8-Hour ozone averages measured at the mobile monitor in comparison to the NAAQS. (Red indicates data from the Battlement Mesa location, Blue indicates Glenwood Springs)




Figure 3-4. Fourth highest and maximum 8-Hour ozone averages measured at the Carbondale site in comparison to the NAAQS.




Figure 3-5. Daily maximum 8-hour averages of ozone monitored at the Rifle site in 2019 in comparison to the NAAQS.




Figure 3-6. Daily maximum 8-hour averages of ozone monitored at the mobile monitor (Battlement Mesa site) in 2019 in comparison to the NAAQS.




Figure 3-7. Daily maximum 8-hour averages of ozone monitored at the Carbondale site in 2019 in comparison to the NAAQS.

3.2 NITROGEN DIOXIDE

Nitrogen Dioxide (NO₂) is generally measured alongside nitric oxide (NO), where NO and NO₂ are the primary components of Oxides of Nitrogen (NO_x). Only NO₂ is considered a regulated pollutant, but all components of NO_x have been identified as precursors for both O₃ and particulate matter. NO is a colorless and odorless gas which, in the presence of O₃, will react to form NO₂. NO₂ is a reddish-brown gas which is partially responsible for the "brown haze" observed near large cities.

The NAAQS for NO₂ include an annual arithmetic mean of 53 ppb and a 1-hour daily maximum of 100 ppb. A violation of the 1-hour standard occurs when the 3-year average of the 98th percentile of the daily maximum 1-hour averages is greater than the standard.

NO₂ is measured at the mobile station, which began at the Battlement Mesa site in late 2012. The mobile site operated at the Glenwood Springs location from February 2015 through February 2016. In March 2016 the mobile monitor returned to the Battlement Mesa monitoring location. Figure 3-8 presents annual average of NO₂ recorded at the mobile monitoring site, and Figure 3-9 presents daily maximum 1-hour averages of NO₂. Plots indicate that measured NO₂ values are well below the NAAQS standards.




Figure 3-8. Annual average nitrogen dioxide monitored at the mobile monitoring station (Battlement Mesa and Glenwood Springs) in comparison to the NAAQS.




Figure 3-9. Daily maximum 1-hour averages of nitrogen dioxide monitored at the Battlement Mesa site in 2019 in comparison to the NAAQS.

3.3 PARTICULATE MATTER

Particulate matter (PM) consists of solid particles and liquid droplets that are small enough to be inhaled. Particles smaller than 10 micrometers in diameter (PM₁₀), are approximately 1/7th the width of a human hair, and pose a health concern because they can be inhaled into and accumulate in the respiratory system. Particles less than 2.5 micrometers in diameter (PM_{2.5}) are referred to as "fine" particles and are believed to pose the greatest health risks. Because of their small size, approximately 1/30th the average width of a human hair, fine particles can lodge deeply into the lungs. Particulate matter can be emitted directly into the air or can be formed in the atmosphere through complex chemical reactions from emissions of sulfur dioxides, nitrogen oxides, and other compounds. Larger particulate matter (PM₁₀) can come from sources such as road dust, construction, and crushing or grinding operations. Fine particulate matter (PM_{2.5}) can come from all kinds of combustion, including motor vehicles, power plants, residential wood burning, and forest fires; as well as from some industrial processes that may contain organic compounds and metals. PM_{2.5} particles are the main cause of reduced visibility or haze. Particulate sources associated with natural gas development may include grading and leveling of well pads, construction of facilities, construction of access roads to well pads, and subsequent vehicle traffic. Naturally occurring emissions such as forest fires can also contribute to particulate matter.

Continuous PM₁₀ and PM_{2.5} monitoring began at the Rifle site in September 2008. The Rifle site stopped collecting continuous PM₁₀ measurements at the end of 2014. Continuous PM₁₀ and PM_{2.5} instruments operated at the mobile monitor beginning in November 2012 and continuous PM_{2.5} monitoring began at the Carbondale site in March 2012. Filter-based 24-hour PM₁₀ monitoring conducted by CDPHE was discontinued as of December 31, 2018. Summaries for both continuous PM₁₀ and PM_{2.5} measurements are provided below.

3.3.1 PM₁₀ Measurements

Figure 3-10 presents the highest and second highest 24-hour average values measured at the mobile site. The level of the national primary and secondary ambient air quality standards for PM₁₀ is a 24-hour average concentration of 150 micrograms per cubic meter ($\mu\text{g}/\text{m}^3$). A violation of the standard occurs when the average number of days with a 24-hour average concentration above 150 $\mu\text{g}/\text{m}^3$ over a 3-year period is greater than one.




Figure 3-10. Highest and second highest 24-hour average PM₁₀ measured at the Mobile Monitor (Battlement Mesa and Glenwood Springs in 2015) site in comparison to the NAAQS.

3.3.2 PM_{2.5} Measurements

Continuous PM_{2.5} data have been monitored at the Rifle site since mid-2008. Continuous PM₁₀ and PM_{2.5} data have been monitored at the Mobile Monitor (Battlement Mesa and Glenwood Springs) site since November 2012, and continuous PM_{2.5} data have been monitored at the Carbondale site since March 2012.

Figure 3-11 presents the annual average of continuous PM_{2.5} measured at all Garfield County sites since 2008. Figures 3-12 through 3-14 present the highest and 98th percentile 24-hour average values measured at the sites. The NAAQS for PM_{2.5} is an arithmetic mean of 12 µg/m³ annually and a 24-hour average of 35 µg/m³. A violation of the PM_{2.5} standard occurs when the 3-year average of the weighted annual mean exceeds that annual standard, or the 3-year average of the 98th percentile 24-hour average value exceeds the 24-hour standard. The highest 24-hour PM_{2.5} values measured in 2008, 2009, and 2012 at the Rifle site were above the standard, but these are not considered violations because the 98th percentile values averaged over a 3-year period were below the standard.




Figure 3-11. Annual average PM_{2.5} measured at the Garfield County sites in comparison to the NAAQS.




Figure 3-12. Highest and 98th percentile 24-hour average PM_{2.5} measured at the Carbondale site in comparison to the NAAQS.




Figure 3-13. Highest and 98th percentile 24-hour average PM_{2.5} measured at the Rifle site. Note that exceedances of the NAAQS are not considered a violation until the 3-year average of the 98th percentile value exceeds the NAAQS.




Figure 3-14. Highest and 98th percentile 24-hour average $PM_{2.5}$ measured at the Mobile Monitor (Battlement Mesa, 2013-2014, 2016-2019, and Glenwood Springs, 2015, sites) in comparison to the NAAQS.

4.0 VOLATILE ORGANIC COMPOUNDS

Volatile organic compounds (VOCs) include some hazardous air pollutants (HAPs), polycyclic aromatic hydrocarbons (PAHs), heavy metals, speciated non-methane organic compounds, and carbonyl compounds; which are known or believed to cause human health effects. Methane, SNMOCs, and carbonyl compounds are subsets of volatile organic compounds (VOCs), which are carbon-based and hydrogen-based chemicals that exist in the gas phase or can evaporate from liquids. VOCs can react in the atmosphere to form ozone (O₃) and particulate matter.

Garfield County has measured speciated non-methane organic compounds (SNMOCs) and carbonyl compounds using a 1-in-6 and 1-in-12 day (respectively) schedule since 2008. Additionally, the County added continuous measurements of methane (CH₄), non-methane, and total hydrocarbons as a part of the Mobile Monitor, which began operation at the Battlement Mesa site in late 2012. The Mobile Monitor operated in Glenwood Springs from February 2015 through February 2016 (data for this location were summarized in the 2015 Annual Data Report). The Mobile Monitor returned to the Battlement Mesa site in March 2016. Summaries of methane, SNMOCs, and carbonyls levels measured in 2019 are presented in this section.

4.1 Continuous Methane and Non-Methane Hydrocarbons

Continuous methane (CH₄) and non-methane organic compounds (NMOC) measurements are collected along with the canister based SNMOC measurements at the mobile monitor. Figure 4-1 presents daily maximum 1-hour concentrations of methane measured at the site in units of ppm. Methane is a pollutant that persists in the atmosphere for long periods of time (~12 years), so there is generally a background concentration of methane present globally even in very remote locations. Methane is also the primary component in natural gas, and can be elevated in oil and gas development areas. Methane is not considered a hazardous air pollutant, but it is a major greenhouse gas and can contribute to climate change. The plot shows a baseline slightly above the global background methane reported by the EPA (~1.8 ppm, see https://19january2017snapshot.epa.gov/climatechange_.html) during the 2019 monitoring period.




Figure 4-1. Daily average methane measured at the Mobile Monitor (Battlement Mesa site) in 2019.

Concentrations of speciated non-methane organic compounds (SNMOC) compounds are available from the canister sample analysis and continuous measurements of total NMOCs are collected at the Mobile Monitoring station. Continuous total NMOC measurements are useful along with canister measurements because continuous measurements are available on an hourly basis in near real-time, while canister samples are only available every sixth day as 24-hour averages. Figure 4-2 presents a time series plot of 24-hour averages of total NMOCs from both continuous and canister based methods, showing good agreement for most of the year.




Figure 4-2. Daily average canister samples and continuous measurements of non-methane organic compounds measured at the Mobile Monitor (Battlement Mesa site) in 2019.

4.2 Speciated Non-Methane Organic Compounds

Garfield County began collecting speciated non-methane organic compounds (SNMOC) data at the Parachute (PACO), Rifle (RECO), and Bell Ranch (BRCO) sites in 2008, at the Mobile Monitor (Battlement Mesa (BMCO) and Glenwood Springs (GSCO)) in September 2010, and at the Carbondale (RFCO) site in 2012. SNMOC compounds are collected and analyzed according to EPA Compendium Method TO-12, with 24-hour samples collected on a 1-in-6 day schedule at four sites, and 1-in-12 in Carbondale. Annual averages are presented here, and Appendix B lists minimum, maximum, and average concentrations of all detected SNMOC compounds by site.

SNMOC compounds can be grouped into classifications with similar characteristics. For summary purposes, measured SNMOC compounds are grouped into the following categories:

- **Light Alkanes:** Alkanes are the simplest hydrocarbons, consisting of only carbon and hydrogen with single bonds. Light alkanes, which include alkanes with up to five carbon atoms (ethane, propane, iso/n-butane and iso/n-pentane), along with methane, are primary components of natural gas and gasoline vapors.
- **Heavy Alkanes:** The hydrocarbons in crude oil are mostly heavy alkanes, which here include alkanes with more than five carbon atoms (C5). Crude oil products include gasoline, a refined mix of predominantly C6 to C10 hydrocarbons, and diesel, which is a refined mix ranging from approximately C10 to C15.
- **Alkenes:** Alkenes are more complex than alkanes, with at least one carbon to carbon double bond. These compounds are not generally found in crude oil. Alkenes are much more reactive than alkanes, and will deplete quickly in the atmosphere. Alkenes are produced in refineries when larger alkane molecules are dissociated (or cracked) into smaller compounds. Some alkene compounds, including terpenes such as isoprene and a- and b-pinene, are naturally emitted from vegetation.
- **Aromatics:** Aromatic compounds are the most abundant compounds emitted from gas-fired engines. These compounds include the BTEX parameters (benzene, toluene, ethylbenzene, and m/p-xylenes), which are commonly associated with motor vehicles, and other engine sources such as those associated with oil and gas production.

Figure 4-3 presents comparisons of annual average SNMOC data collected between 2008 and 2019, showing that total SNMOC measurements were generally decreasing since 2008, but showed some increases in 2019 at the Bell Ranch and Carbondale sites. Figure 4-4 presents categories of measured SNMOCs in units of ppbV (parts per billion by volume) measured in 2019 at each site. In general, measured compounds consisted mostly of light alkanes. Light alkanes represented between 68% and 88% of total SNMOCs measured at sites near natural gas development. These trends can be influenced by the variations in temperature, as VOCs can be trapped under inversion layers during the colder months, especially if wind conditions are stagnant. Also, some emissions, including cold-start engine emissions and residential wood burning, are higher in the winter.

Figure 4-5 presents measurements by category in units of ppbC, where ppbC represents the number of carbon molecules measured (ppbV multiplied by the number of carbons in each compound). Heavier alkanes and aromatics are more significant sources of carbon than the lighter alkanes. The unknown category indicates the part of the total carbon measurements where individual species were not identified.

Carbon content in a molecule is important because it is related to compound reactivity, which contributes to O₃ formation potential. O₃ is formed from photochemical interactions of VOCs and NO_x in the presence of sunlight, as described in Section 3.1. The light alkanes that dominate measurements by volume are the least reactive compounds but could theoretically contribute significantly to O₃ formation potential. Highly reactive compounds including aromatics such as toluene and m/p-xylenes are less abundant, but have greater potential to contribute to the O₃ formation due to their higher reactivity. Currently, Garfield County is not in violation of the NAAQS for O₃. However, if O₃ levels become more of a concern in Garfield County, it would be useful to target further controls for emissions of VOCs that have the greatest potential to contribute to O₃ formation.




Figure 4-3 Average SNMOC concentrations measured at Garfield County sites between 2008 and 2019.




Figure 4-4. 2019 24-hour SNMOC measurements by category in units of ppbV.




Figure 4-5. 2019 24-hour SNMOC measurements by category in units of ppbC.

4.3 CARBONYLS

Garfield County began collecting carbonyl data at the Parachute, Rifle, and Bell Ranch sites in 2008, and at the mobile monitor (Battlement Mesa (BMCO) and Glenwood Springs (GSCO)) site in September 2010. The Carbondale (RFCO) site collected carbonyl samples from June 2012 through December 2014. These compounds were collected and analyzed according to EPA Compendium Method TO-11A, with 24-hour samples collected at all four sites on a 1-in-12 day schedule. This method includes analysis for 12 different carbonyl compounds. Carbonyls are highly reactive and play a critical role in the formation of O₃. Some carbonyls, including formaldehyde and acetaldehyde, also have adverse chronic and acute health effects. The major sources of directly emitted carbonyls are fuel combustion, mobile sources, and process emissions from oil refineries.

Figure 4-6 presents comparisons of annual average carbonyl data collected between 2008 and 2019. Figure 4-7 presents time series plots of the major compounds, and Appendix C lists minimum, maximum, and average concentrations of all detected carbonyl compounds. Major compounds measured included formaldehyde, acetaldehyde, and acetone.




Figure 4-6. Average carbonyl concentrations measured at Garfield County sites between 2008 and 2019.




Figure 4-7. 2019 24-hour major carbonyl compound concentrations in units of ppbV.

4.4 HAZARDOUS AIR POLLUTANTS (HAPS)

The EPA has designated approximately 190 compounds as HAPs, including benzene, toluene, ethyl-benzene and xylenes (also known as the BTEX compounds). No NAAQS or any other ambient air standards exist for HAPs. Instead, emissions limits on industrial sources have been set, and the EPA has developed a set of risk factors for both acute and chronic exposures for HAPs. In addition, risk factors from the Agency for Toxic Substances and Disease Registry (ATSDR), the California Air Resources Board (CARB), the National Institute for Occupational Safety and Health (NIOSH), and others can be used to determine potential risks from exposure to VOCs.

Of the 78 SNMOC and 12 carbonyl compounds measured in Garfield County, 21 compounds are considered HAPs. *The Garfield County Air Toxics Inhalation Screening Level Human Risk Assessment* (CDPHE 2010) assessed data collected in 2008. Risk assessments based on 2008 through 2012 VOC levels were assessed in a separate risk assessment report prepared by the CDPHE Disease Control and Environmental Epidemiology Division in 2015; available on the Garfield County Air Quality Management website (<http://www.garfield-county.com/air-quality/2008-12-health-risk-assessment.aspx>). Findings of the 2010 report indicated that, individually, the HAP components were below risk assessment criteria, but cumulative effects approached chronic (70 year exposure period) non-hazard levels. The largest contributors to the cumulative levels were benzene and formaldehyde. Findings of the 2015 report indicate a trend toward decreasing cumulative cancer risk estimates and noncancer hazards for all contaminants of potential concerns except 1,2-butadiene at all monitoring sites. Definitive conclusions cannot be drawn regarding the magnitude of cancer and noncancer risks associated with emissions from oil and gas operations because of the major uncertainties and knowledge gaps.

Summaries in this report include annual averages for the HAPs measured in Garfield County, but do not address health effects of these compounds. It is important to note that annual average values summarized here do not necessarily indicate a health risk, as actual magnitudes of these HAP compounds related to possible health risk are evaluated separately in the CDPHE risk assessment reports.

4.4.1 Annual Average HAP Concentrations

Annual trends were calculated for each HAP, with a trend defined as the slope derived from Theil statistics. This is a nonparametric regression technique commonly applied to environmental data to determine statistically significant trends. The significance of the trend is represented with p-values calculated using Mann-Kendall trend statistics. Determining a significance level helps to distinguish random variability in data from a real tendency to increase or decrease over time, where lower p-values indicate higher confidence levels in the computed slopes.

Tables 4-1 through 4-5 present annual averages and trends for HAP concentrations measured between 2008 and 2019. For purposes here, regional trends are considered statistically significant if the p-value statistic is less than 0.10 (90% confidence level). Statistically significant decreasing trends are indicated in blue, and statistically significant increasing trends

are depicted in red. Data collected since 2008 indicate that a number of HAPs compounds have measured statistically significant decreasing annual average trends at all sites.

Table 4-1
Parachute Site
Annual Average Mass Trends (HAPs Parameters) 2008-2019

Parameter	Average Mass ($\mu\text{g}/\text{m}^3$)												Slope ($\mu\text{g}/\text{m}^3$ per year)	p- Value
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019		
1,2,4-Trimethylbenzene	0.91	0.62	0.42	0.43	0.38	0.79	0.50	0.36	0.53	0.57	0.39	0.30	-0.02	0.06
1,3,5-Trimethylbenzene	0.61	0.51	0.40	0.22	0.17	0.32	0.24	0.16	0.20	0.21	0.22	0.12	-0.03	0.00
1,3-Butadiene	0.08	0.12	0.10	0.10	0.09	0.06	0.07	0.03	0.03	0.02	0.04	0.02	-0.01	0.00
Acetaldehyde	1.11	0.99	0.92	0.90	0.69	0.73	0.48	0.33	0.59	0.60	0.82	0.65	-0.05	0.01
Acetone	3.42	3.28	2.67	2.79	2.28	2.29	1.60	1.45	2.27	2.38	3.17	2.81	-0.06	0.19
Benzene	2.31	2.69	1.74	1.44	1.25	1.85	1.49	1.21	1.20	1.01	1.09	0.94	-0.12	0.00
Crotonaldehyde	0.10	0.10	0.12	0.09	0.06	0.07	0.04	0.05	0.06	0.08	0.09	0.06	0.00	0.06
Cyclohexane	3.92	3.77	2.82	2.22	2.00	3.15	2.18	1.73	1.35	1.39	1.63	1.31	-0.22	0.00
Ethylbenzene	0.59	0.44	1.02	0.32	0.16	0.23	0.19	0.16	0.17	0.39	0.97	0.16	-0.02	0.06
Formaldehyde	1.74	1.73	1.53	1.64	1.20	1.28	0.97	0.84	1.18	1.21	1.58	1.29	-0.04	0.08
Isopropylbenzene	0.09	0.08	0.07	0.08	0.06	0.06	0.06	0.04	0.09	0.07	0.05	0.04	0.00	0.02
Methylcyclohexane	9.24	9.43	6.23	4.65	4.18	6.59	4.62	3.60	2.71	3.02	3.71	2.78	-0.53	0.00
m-Xylene/p-Xylene	3.91	3.63	2.19	1.11	1.13	1.80	1.48	1.08	0.90	0.93	1.70	0.78	-0.18	0.00
n-Hexane	5.78	5.64	3.93	3.34	3.04	3.12	3.06	2.48	1.99	1.83	2.21	1.95	-0.29	0.00
n-Nonane	2.20	2.01	1.13	0.97	0.59	0.87	0.69	0.47	0.40	0.54	0.78	0.39	-0.12	0.00
n-Propylbenzene	0.18	0.15	0.13	0.10	0.10	0.15	0.13	0.06	0.12	0.11	0.08	0.08	-0.01	0.01
o-Xylene	0.77	0.65	0.43	0.40	0.26	0.40	0.33	0.25	0.25	0.49	0.34	0.18	-0.03	0.00
Propionaldehyde	0.12	0.09	0.09	0.09	0.07	0.07	0.04	0.03	0.06	0.07	0.07	0.07	0.00	0.02
Propylene	0.57	0.57	0.62	0.55	0.70	0.49	0.63	0.45	0.51	0.44	0.45	0.55	-0.01	0.04
Styrene	0.12	0.08	0.12	0.13	1.41	3.76	0.92	0.82	1.31	1.62	0.96	0.11	0.09	0.10
Toluene	9.86	5.83	3.94	5.79	4.18	4.45	4.54	2.86	9.61	3.94	4.62	1.92	-0.25	0.06

Table 4-2
Rifle Site
Annual Average Mass Trends (HAPs Parameters) 2008-2019

Parameter	Average Mass ($\mu\text{g}/\text{m}^3$)												Slope ($\mu\text{g}/\text{m}^3$ per year)	p- Value
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019		
1,2,4-Trimethylbenzene	0.62	0.71	0.52	0.42	0.49	0.76	0.77	0.58	0.70	0.62	0.51	0.47	-0.01	0.27
1,3,5-Trimethylbenzene	0.33	0.33	0.30	0.18	0.16	0.25	0.24	0.20	0.21	0.17	0.14	0.13	-0.02	0.00
1,3-Butadiene	0.13	0.12	0.17	0.19	0.18	0.12	0.12	0.09	0.09	0.06	0.07	0.06	-0.01	0.00
Acetaldehyde	1.55	1.39	1.47	1.41	1.04	0.92	0.51	0.77	0.86	0.99	0.98	0.79	-0.06	0.01
Acetone	3.58	3.11	3.08	2.74	2.62	2.24	1.51	2.44	2.54	2.80	2.99	2.45	-0.06	0.03
Benzene	1.68	2.22	1.44	1.27	0.99	1.52	1.09	1.01	1.11	0.92	0.82	0.80	-0.08	0.00
Crotonaldehyde	0.15	0.17	0.18	0.14	0.10	0.09	0.07	0.09	0.08	0.12	0.10	0.09	-0.01	0.01
Cyclohexane	2.46	2.48	1.84	1.86	1.64	2.88	1.74	1.30	1.44	0.89	0.79	0.73	-0.15	0.00
Ethylbenzene	0.48	0.56	0.84	0.35	0.26	0.35	0.32	0.31	0.29	0.49	0.21	0.18	-0.03	0.01
Formaldehyde	1.89	1.67	1.80	1.82	1.39	1.38	0.74	1.14	1.31	1.47	1.34	1.17	-0.05	0.01
Isopropylbenzene	0.08	0.08	0.08	0.08	0.06	0.06	0.06	0.05	0.05	0.07	0.04	0.04	0.00	0.00
Methylcyclohexane	4.78	5.08	3.55	3.36	2.70	4.63	2.67	1.88	1.74	1.18	1.09	1.06	-0.34	0.00
m-Xylene/p-Xylene	2.35	2.58	1.70	0.90	1.07	1.59	1.28	1.25	1.06	0.79	0.80	0.69	-0.14	0.00
n-Hexane	4.50	4.61	3.48	4.40	3.05	3.41	3.12	2.61	2.19	1.64	1.44	1.51	-0.31	0.00
n-Nonane	0.81	0.86	0.59	0.44	0.31	0.48	0.34	0.27	0.36	0.18	0.19	0.18	-0.06	0.00
n-Propylbenzene	0.15	0.16	0.16	0.12	0.12	0.14	0.16	0.10	0.12	0.12	0.10	0.09	-0.01	0.00
o-Xylene	0.64	0.75	0.51	0.43	0.36	0.49	0.44	0.46	0.40	0.58	0.31	0.26	-0.03	0.01
Propionaldehyde	0.16	0.15	0.16	0.13	0.10	0.11	0.06	0.10	0.10	0.14	0.12	0.07	-0.01	0.02
Propylene	0.86	1.05	1.04	0.93	0.98	0.83	0.95	0.81	0.82	0.69	0.69	0.71	-0.03	0.00
Styrene	0.08	0.08	0.08	0.10	0.68	2.13	2.18	0.51	0.97	1.47	0.44	0.20	0.05	0.06
Toluene	4.34	4.62	3.07	5.90	2.16	3.25	2.79	3.46	2.27	1.97	1.81	1.56	-0.26	0.00

Table 4-3
Bell Ranch Site
Annual Average Mass Trends (HAPs Parameters) 2008-2019

Parameter	Average Mass ($\mu\text{g}/\text{m}^3$)												Slope ($\mu\text{g}/\text{m}^3$ per year)	p- Value
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019		
1,2,4-Trimethylbenzene	0.28	0.28	0.22	0.21	0.21	0.38	0.31	0.17	0.26	0.25	0.15	0.16	-0.01	0.04
1,3,5-Trimethylbenzene	0.16	0.15	0.15	0.10	0.07	0.12	0.10	0.07	0.07	0.07	0.06	0.04	-0.01	0.00
1,3-Butadiene	0.05	0.06	0.06	0.06	0.10	0.06	0.06	0.03	0.04	0.02	0.03	0.02	0.00	0.03
Acetaldehyde	0.83	0.79	0.84	0.61	0.61	0.56	0.51	0.47	0.22	0.61	0.82	0.33	-0.04	0.01
Acetone	2.77	2.66	2.50	2.35	2.28	1.99	2.03	2.00	1.01	2.50	3.21	1.38	-0.11	0.06
Benzene	1.34	1.39	1.08	0.86	0.67	1.14	0.70	0.65	0.69	0.46	0.43	0.44	-0.08	0.00
Crotonaldehyde	0.11	0.15	0.15	0.04	0.09	0.11	0.07	0.09	0.04	0.12	0.20	0.06	0.00	0.37
Cyclohexane	5.08	2.73	2.32	1.99	1.77	2.66	1.71	1.32	1.22	0.79	0.83	0.75	-0.21	0.00
Ethylbenzene	0.26	0.24	0.36	0.16	0.08	0.12	0.09	0.06	0.06	0.30	0.06	0.04	-0.02	0.00
Formaldehyde	1.00	1.37	1.20	0.96	1.02	0.87	0.80	0.76	0.37	1.02	1.36	0.56	-0.05	0.06
Isopropylbenzene	0.08	0.07	0.07	0.07	0.05	0.05	0.05	0.04	0.03	0.05	0.02	0.03	0.00	0.00
Methylcyclohexane	5.87	4.99	4.14	3.63	3.15	4.59	2.85	2.25	1.98	1.30	1.51	1.20	-0.39	0.00
m-Xylene/p-Xylene	1.36	1.28	0.98	0.45	0.50	0.78	0.46	0.33	0.33	0.22	0.31	0.20	-0.10	0.00
n-Hexane	6.39	5.35	4.40	3.46	3.33	3.22	3.04	2.43	2.18	1.30	1.39	1.58	-0.39	0.00
n-Nonane	0.68	0.62	0.46	0.33	0.25	0.37	0.22	0.16	0.16	0.11	0.19	0.12	-0.05	0.00
n-Propylbenzene	0.08	0.08	0.09	0.08	0.07	0.09	0.11	0.04	0.05	0.07	0.04	0.04	0.00	0.02
o-Xylene	0.31	0.30	0.23	0.18	0.13	0.20	0.23	0.16	0.12	0.35	0.10	0.07	-0.02	0.01
Propionaldehyde	0.08	0.09	0.09	0.06	0.07	0.06	0.06	0.06	0.03	0.10	0.11	0.06	0.00	0.23
Propylene	0.27	0.42	0.49	0.38	0.38	0.39	0.53	0.30	0.38	0.32	0.29	0.38	0.00	0.16
Styrene	0.12	0.08	0.08	0.14	0.88	1.52	0.69	0.22	0.95	0.23	0.08	0.07	0.00	0.53
Toluene	3.63	2.52	1.90	1.53	1.39	3.56	2.13	2.37	1.84	0.76	0.93	0.77	-0.18	0.01

Table 4-4
Mobile Monitor – Battlement Mesa Site
Annual Average Mass Trends (HAPs Parameters) 2008-2019

Parameter	Average Mass ($\mu\text{g}/\text{m}^3$)												Slope ($\mu\text{g}/\text{m}^3$ per year)	p- Value
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019		
1,2,4-Trimethylbenzene			0.32	0.44	0.35	0.40	0.26	--	0.23	0.37	0.22	0.18	-0.02	0.02
1,3,5-Trimethylbenzene			0.24	0.22	0.15	0.16	0.14	--	0.14	0.16	0.14	0.06	-0.01	0.01
1,3-Butadiene			0.06	0.06	0.10	0.05	0.05	--	0.03	0.02	0.03	0.02	-0.01	0.00
Acetaldehyde			0.89	0.76	0.53	0.44	0.41	--	0.48	0.44	0.75	0.63	-0.02	0.19
Acetone			5.34	3.83	2.86	1.76	1.70	--	2.06	2.01	2.81	2.64	-0.15	0.15
Benzene			1.48	1.56	1.09	1.24	0.88	--	0.79	0.85	0.66	0.48	-0.11	0.00
Crotonaldehyde			0.07	0.07	0.05	0.06	0.06	--	0.08	0.07	0.11	0.10	0.00	0.15
Cyclohexane			2.51	2.09	1.77	2.40	1.74	--	1.49	1.57	1.36	0.80	-0.14	0.00
Ethylbenzene			0.28	0.37	0.19	0.14	0.11	--	0.09	0.35	0.14	0.08	-0.02	0.01
Formaldehyde			1.38	1.45	0.87	0.82	0.75	--	0.93	0.88	1.38	1.20	-0.01	0.43
Isopropylbenzene			0.06	0.08	0.05	0.05	0.05	--	0.03	0.04	0.02	0.04	0.00	0.00
Methylcyclohexane			5.13	4.23	3.57	5.06	3.64	--	3.37	3.69	3.40	1.78	-0.22	0.01
m-Xylene/p-Xylene			1.44	1.12	1.06	1.23	0.94	--	0.82	0.91	0.98	0.42	-0.06	0.00
n-Hexane			4.02	3.34	2.89	2.42	2.35	--	1.91	1.95	1.73	1.31	-0.23	0.00
n-Nonane			0.59	0.68	0.43	0.62	0.48	--	0.48	0.57	0.64	0.23	-0.02	0.19
n-Propylbenzene			0.10	0.14	0.09	0.05	0.09	--	0.03	0.08	0.05	0.05	-0.01	0.02
o-Xylene			0.35	0.46	0.27	0.26	0.19	--	0.18	0.45	0.21	0.10	-0.03	0.02
Propionaldehyde			0.10	0.08	0.06	0.05	0.04	--	0.04	0.06	0.08	0.07	0.00	0.30
Propylene			0.51	0.42	0.44	0.36	0.40	--	0.33	0.33	0.32	0.42	-0.01	0.04
Styrene			0.07	0.08	1.54	0.18	0.15	--	0.24	0.27	0.07	0.01	0.00	0.50
Toluene			2.87	2.99	4.07	5.58	2.54	--	2.63	2.68	2.19	1.21	-0.18	0.02

(--) No data available.

Table 4-5
Carbondale Site
Annual Average Mass Trends (HAPs Parameters) 2008-2019

Parameter	Average Mass ($\mu\text{g}/\text{m}^3$)												Slope ($\mu\text{g}/\text{m}^3$ per year)	p- Value
	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019		
1,2,4-Trimethylbenzene	--	--	--	--	0.33	0.59	0.60	0.48	0.38	0.32	0.24	0.25	-0.06	0.03
1,3,5-Trimethylbenzene	--	--	--	--	0.09	0.13	0.13	0.07	0.08	0.09	0.05	0.05	-0.01	0.05
1,3-Butadiene	--	--	--	--	0.14	0.06	0.06	0.04	0.04	0.02	0.03	0.03	-0.01	0.00
Acetaldehyde	--	--	--	--	0.57	0.58	0.60	--	--	--	--	--		
Acetone	--	--	--	--	1.67	1.62	2.01	--	--	--	--	--		
Benzene	--	--	--	--	0.46	0.47	0.45	0.85	0.45	0.38	0.35	0.34	-0.02	0.02
Crotonaldehyde	--	--	--	--	0.15	0.12	0.11							
Cyclohexane	--	--	--	--	0.34	0.42	0.23	0.18	0.18	0.18	0.15	0.18	-0.02	0.01
Ethylbenzene	--	--	--	--	0.10	0.12	0.11	0.12	0.10	0.34	0.10	0.09	0.00	0.14
Formaldehyde	--	--	--	--	0.83	0.75	0.79	--	--	--	--	--		
Isopropylbenzene	--	--	--	--	0.05	0.06	0.06	0.04	0.04	0.05	0.02	0.04	0.00	0.05
Methylcyclohexane	--	--	--	--	0.50	0.67	0.31	0.17	0.16	0.17	0.15	0.13	-0.05	0.00
m-Xylene/p-Xylene	--	--	--	--	0.31	0.49	0.41	0.41	0.34	0.28	0.28	0.24	-0.03	0.01
n-Hexane	--	--	--	--	0.84	0.65	0.58	0.47	0.48	0.39	0.39	0.51	-0.05	0.03
n-Nonane	--	--	--	--	0.10	0.13	0.16	0.13	0.13	0.09	0.08	0.08	-0.01	0.05
n-Propylbenzene	--	--	--	--	0.11	0.14	0.19	0.08	0.05	0.09	0.05	0.05	-0.01	0.03
o-Xylene	--	--	--	--	0.14	0.21	0.20	0.21	0.17	0.40	0.13	0.11	-0.01	0.20
Propionaldehyde	--	--	--	--	0.08	0.07	0.08	--	--	--	--	--		
Propylene	--	--	--	--	0.49	0.40	0.91	0.63	0.47	0.40	0.36	0.42	-0.02	0.09
Styrene	--	--	--	--	1.27	2.80	0.77	0.36	0.22	0.45	0.10	0.14	-0.17	0.01
Toluene	--	--	--	--	3.86	4.37	1.89	17.05	1.96	1.09	1.27	1.60	-0.39	0.09

(P) Partial Year (--) No data available.

5.0 REFERENCES

- Colorado Department of Public Health and Environment (CDPHE). 2010. *Garfield County Air Toxics Inhalation: Screening Level Human Health Risk Assessment*. Available online at <http://www.garfield-county.com/public-health/documents/>.
- Colorado Department of Public Health and Environment (CDPHE). 2015. *Garfield County Air Toxics Inhalation: Screening Level Human Health Risk Assessment: Trends in Air Quality from 2008 to 2012*. Available online at <http://www.garfield-county.com/air-quality/2008-12-health-risk-assessment.aspx/>.
- Environmental Protection Agency (EPA). 2011. *2008-2009 National Monitoring Programs Annual Report (UATMP, NATTS, and CSATAM)*. Available online at <http://www.epa.gov/ttnamti1/uatm.html>.

APPENDIX A

Quarterly Time Series Plots

Acronyms Used on Plots:

TNMOC = Total Non-Methane Organic Carbon

CH₄ = Methane

NMHC = Continuous Non-Methane Hydrocarbons

O₃ = Ozone

NO = Nitric Oxide

NO₂ = Nitrogen Dioxide

NO_x = Oxides of Nitrogen

PM₁₀ = Particulate Matter ≤ 10 μm

PM_{2.5} = Particulate Matter ≤ 2.5 μm


RH = Relative Humidity

AT = Atmospheric Temperature


VWD = Vector Wind Direction

SWS = Scalar Wind Speed


Garfield County, CO
Parachute Site




Garfield County, CO
Parachute Site




Garfield County, CO
Parachute Site




Garfield County, CO
Parachute Site




Garfield County, CO
Rifle Site




Garfield County, CO
Rifle Site




Garfield County, CO
Rifle Site




Garfield County, CO
Rifle Site




Garfield County, CO
Battlement Mesa Site




Garfield County, CO
Battlement Mesa Site




Garfield County, CO
Battlement Mesa Site




Garfield County, CO
Battlement Mesa Site




Garfield County, CO
Bell Ranch Site




Garfield County, CO
Bell Ranch Site




Garfield County, CO
Bell Ranch Site




Garfield County, CO
Bell Ranch Site




Garfield County, CO
Carbondale Site




Garfield County, CO
Carbondale Site



Garfield County, CO
Carbondale Site



Garfield County, CO
Carbondale Site



APPENDIX B
SNMOC Concentrations

Garfield County SNMOC Monitoring
Parachute (PACO)
1/3/2019-11/17/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
1,2,3-Trimethylbenzene (526-73-8)	34	11	0.01	0.06	0.01
1,2,4-Trimethylbenzene (95-63-6)	34	31	0.01	0.33	0.06
1,3,5-Trimethylbenzene (108-67-8)	34	25	0.01	0.06	0.03
1,3-Butadiene (106-99-0)	34	8	0.00	0.03	0.01
1-Dodecene (112-41-4)	34	6	0.01	0.16	0.03
1-Heptene (592-76-7)	34	1	0.03	0.03	0.01
1-Hexene (592-41-6)	34	22	0.01	0.05	0.01
1-Nonene (124-11-8)	34	26	0.01	0.10	0.03
1-Octene (111-66-0)	34	34	0.01	0.16	0.05
1-Pentene (109-67-1)	34	34	0.01	0.12	0.04
1-Undecene (821-95-4)	34	2	0.01	0.04	0.03
2,2,3-Trimethylpentane (564-02-3)	34	13	0.02	0.05	0.01
2,2-Dimethylbutane (75-83-2)	34	31	0.03	0.20	0.06
2,3,4-Trimethylpentane (565-75-3)	34	15	0.01	0.06	0.01
2,3-Dimethylbutane (79-29-8)	34	34	0.04	0.33	0.11
2,3-Dimethylpentane (565-59-3)	34	34	0.02	0.14	0.05
2,4-Dimethylpentane (108-08-7)	34	34	0.01	0.11	0.04
2-Methyl-1-butene (563-46-2)	34	15	0.01	0.05	0.02
2-Methyl-2-butene (513-35-9)	34	13	0.01	0.07	0.02
2-Methylheptane (592-27-8)	34	33	0.03	0.22	0.08
2-Methylhexane (591-76-4)	34	21	0.07	0.50	0.14
2-Methylpentane (107-83-5)	34	32	0.16	1.12	0.45
3-Methyl-1-butene (563-45-1)	34	1	0.02	0.02	0.00
3-Methylheptane (589-81-1)	34	34	0.02	0.19	0.06
3-Methylhexane (589-34-4)	34	32	0.06	0.45	0.17
3-Methylpentane (96-14-0)	34	34	0.08	0.67	0.24
4-Methyl-1-pentene (691-37-2)	34	1	0.02	0.02	0.01
Acetylene (74-86-2)	34	34	0.14	1.44	0.44
a-Pinene (80-56-8)	34	8	0.01	0.04	0.01
Benzene (71-43-2)	34	34	0.13	0.70	0.29
cis-2-Butene (590-18-1)	34	26	0.01	0.05	0.02
cis-2-Hexene (7688-21-3)	34	2	0.02	0.02	0.01
cis-2-Pentene (627-20-3)	34	8	0.01	0.01	0.00
Cyclohexane (110-82-7)	34	34	0.14	1.08	0.38
Cyclopentane (287-92-3)	34	34	0.03	0.19	0.07
Cyclopentene (142-29-0)	34	8	0.03	0.21	0.05
Ethane (74-84-0)	34	34	12.15	137.00	36.31
Ethylbenzene (100-41-4)	34	33	0.01	0.34	0.04
Ethylene (74-85-1)	34	34	0.40	2.31	1.23
Isobutane (75-28-5)	34	34	0.80	9.38	2.66
Isopentane (78-78-4)	34	3	0.48	4.74	0.19
Isoprene (78-79-5)	34	21	0.01	0.40	0.09

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Parachute (PACO)
1/3/2019-11/17/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
Isopropylbenzene (98-82-8)	34	4	0.01	0.06	0.01
m-Diethylbenzene (141-93-5)	34	5	0.02	0.15	0.02
Methylcyclohexane (108-87-2)	34	34	0.26	1.97	0.69
Methylcyclopentane (96-37-7)	34	33	0.12	0.88	0.32
m-Ethyltoluene (620-14-4)	34	27	0.01	0.20	0.04
m-Xylene/p-Xylene (108-38-3/106-42-3)	34	33	0.06	0.46	0.18
n-Butane (106-97-8)	34	34	0.89	8.93	2.77
n-Decane (124-18-5)	34	28	0.02	0.18	0.05
n-Dodecane (112-40-3)	34	7	0.01	0.04	0.01
n-Heptane (142-82-5)	34	34	0.11	0.77	0.29
n-Hexane (110-54-3)	34	34	0.20	1.48	0.55
n-Nonane (111-84-2)	34	34	0.03	0.20	0.08
n-Octane (111-65-9)	34	34	0.08	0.56	0.20
n-Pentane (109-66-0)	34	34	0.33	2.72	1.01
n-Propylbenzene (103-65-1)	34	10	0.01	0.15	0.02
n-Undecane (1120-21-4)	34	30	0.01	0.06	0.02
o-Ethyltoluene (611-14-3)	34	15	0.01	0.04	0.01
o-Xylene (95-47-6)	34	34	0.01	0.09	0.04
p-Diethylbenzene (105-05-5)	34	2	0.03	0.07	0.01
p-Ethyltoluene (622-96-8)	34	12	0.01	0.11	0.02
Propane (74-98-6)	34	34	3.53	38.67	10.91
Propylene (115-07-1)	34	34	0.09	1.25	0.32
Styrene (100-42-5)	34	2	0.25	0.50	0.02
Toluene (108-88-3)	34	34	0.18	1.05	0.51
trans-2-Butene (624-64-6)	34	15	0.02	0.25	0.04
trans-2-Hexene (4050-45-7)	34	1	0.01	0.01	0.01
trans-2-Pentene (646-04-8)	34	23	0.01	0.06	0.02

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Rifle Library (RECO)
1/3/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
1,2,3-Trimethylbenzene (526-73-8)	46	28	0.01	0.13	0.02
1,2,4-Trimethylbenzene (95-63-6)	46	43	0.02	0.52	0.10
1,3,5-Trimethylbenzene (108-67-8)	46	39	0.01	0.07	0.03
1,3-Butadiene (106-99-0)	46	38	0.01	0.08	0.03
1-Dodecene (112-41-4)	46	6	0.02	0.11	0.03
1-Heptene (592-76-7)	46	40	0.02	0.15	0.06
1-Hexene (592-41-6)	46	27	0.01	0.15	0.02
1-Nonene (124-11-8)	46	41	0.01	0.07	0.02
1-Octene (111-66-0)	46	41	0.01	0.12	0.03
1-Pentene (109-67-1)	46	45	0.02	0.20	0.06
1-Undecene (821-95-4)	46	6	0.01	0.02	0.03
2,2,3-Trimethylpentane (564-02-3)	46	9	0.02	0.04	0.01
2,2,4-Trimethylpentane (540-84-1)	46	40	0.02	0.11	0.05
2,2-Dimethylbutane (75-83-2)	46	42	0.01	0.19	0.04
2,3,4-Trimethylpentane (565-75-3)	46	32	0.01	0.07	0.02
2,3-Dimethylbutane (79-29-8)	46	46	0.02	0.26	0.08
2,3-Dimethylpentane (565-59-3)	46	46	0.03	0.14	0.05
2,4-Dimethylpentane (108-08-7)	46	46	0.02	0.09	0.03
2-Methyl-1-butene (563-46-2)	46	41	0.02	0.16	0.05
2-Methyl-1-pentene (763-29-1)	46	4	0.01	0.01	0.01
2-Methyl-2-butene (513-35-9)	46	39	0.03	0.20	0.07
2-Methylheptane (592-27-8)	46	45	0.01	0.12	0.04
2-Methylhexane (591-76-4)	46	18	0.09	0.42	0.07
2-Methylpentane (107-83-5)	46	40	0.14	1.12	0.40
3-Methylheptane (589-81-1)	46	46	0.02	0.09	0.04
3-Methylhexane (589-34-4)	46	42	0.06	0.34	0.12
3-Methylpentane (96-14-0)	46	44	0.07	0.59	0.19
Acetylene (74-86-2)	46	46	0.31	8.00	1.12
a-Pinene (80-56-8)	46	15	0.00	2.27	0.13
Benzene (71-43-2)	46	46	0.12	0.64	0.25
cis-2-Butene (590-18-1)	46	46	0.02	0.52	0.07
cis-2-Hexene (7688-21-3)	46	4	0.01	0.02	0.01
cis-2-Pentene (627-20-3)	46	42	0.01	0.06	0.02
Cyclohexane (110-82-7)	46	46	0.07	0.61	0.21
Cyclopentane (287-92-3)	46	46	0.02	0.22	0.07
Cyclopentene (142-29-0)	46	8	0.04	0.08	0.05
Ethane (74-84-0)	46	46	5.90	85.00	20.86
Ethylbenzene (100-41-4)	46	45	0.02	0.11	0.04
Ethylene (74-85-1)	46	46	0.65	4.78	1.76
Isobutane (75-28-5)	46	46	0.57	12.00	2.29
Isopentane (78-78-4)	46	2	1.29	5.92	0.16
Isoprene (78-79-5)	46	44	0.01	0.59	0.12

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Rifle Library (RECO)
1/3/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
Isopropylbenzene (98-82-8)	46	13	0.01	0.02	0.01
m-Diethylbenzene (141-93-5)	46	7	0.01	0.16	0.02
Methylcyclohexane (108-87-2)	46	44	0.10	0.80	0.26
Methylcyclopentane (96-37-7)	46	45	0.09	0.60	0.22
m-Ethyltoluene (620-14-4)	46	46	0.02	0.26	0.08
m-Xylene/p-Xylene (108-38-3/106-42-3)	46	46	0.08	0.39	0.16
n-Butane (106-97-8)	46	46	0.85	14.65	3.08
n-Decane (124-18-5)	46	43	0.01	0.17	0.04
n-Dodecane (112-40-3)	46	8	0.02	0.06	0.01
n-Heptane (142-82-5)	46	46	0.07	0.47	0.16
n-Hexane (110-54-3)	46	46	0.20	1.18	0.43
n-Nonane (111-84-2)	46	46	0.01	0.07	0.03
n-Octane (111-65-9)	46	43	0.04	0.22	0.09
n-Pentane (109-66-0)	46	46	0.37	3.74	1.02
n-Propylbenzene (103-65-1)	46	34	0.01	0.09	0.02
n-Undecane (1120-21-4)	46	38	0.01	0.04	0.01
o-Ethyltoluene (611-14-3)	46	36	0.01	0.06	0.02
o-Xylene (95-47-6)	46	46	0.02	0.14	0.06
p-Diethylbenzene (105-05-5)	46	1	0.04	0.04	0.00
p-Ethyltoluene (622-96-8)	46	41	0.01	0.15	0.03
Propane (74-98-6)	46	46	2.08	27.47	7.46
Propylene (115-07-1)	46	46	0.16	1.37	0.41
Propyne (74-99-7)	46	8	0.01	0.06	0.01
Styrene (100-42-5)	46	5	0.06	0.79	0.05
Toluene (108-88-3)	46	46	0.21	0.98	0.42
trans-2-Butene (624-64-6)	46	28	0.02	0.76	0.10
trans-2-Hexene (4050-45-7)	46	6	0.01	0.02	0.01
trans-2-Pentene (646-04-8)	46	45	0.01	0.12	0.04

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Bell Ranch (BRCO)
1/3/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
1,2,3-Trimethylbenzene (526-73-8)	52	10	0.01	0.02	0.00
1,2,4-Trimethylbenzene (95-63-6)	52	37	0.01	0.13	0.03
1,3,5-Trimethylbenzene (108-67-8)	52	10	0.01	0.02	0.01
1,3-Butadiene (106-99-0)	52	5	0.01	0.01	0.01
1-Dodecene (112-41-4)	52	4	0.03	0.28	0.03
1-Heptene (592-76-7)	52	7	0.03	0.05	0.01
1-Hexene (592-41-6)	52	35	0.01	0.06	0.02
1-Nonene (124-11-8)	52	47	0.01	0.06	0.03
1-Octene (111-66-0)	52	52	0.01	0.12	0.04
1-Pentene (109-67-1)	52	49	0.01	0.08	0.04
1-Undecene (821-95-4)	52	13	0.01	0.02	0.03
2,2,3-Trimethylpentane (564-02-3)	52	10	0.01	0.05	0.01
2,2,4-Trimethylpentane (540-84-1)	52	6	0.01	0.03	0.00
2,2-Dimethylbutane (75-83-2)	52	47	0.02	0.11	0.03
2,3,4-Trimethylpentane (565-75-3)	52	19	0.01	0.04	0.01
2,3-Dimethylbutane (79-29-8)	52	52	0.02	0.20	0.06
2,3-Dimethylpentane (565-59-3)	52	51	0.01	0.07	0.03
2,4-Dimethylpentane (108-08-7)	52	48	0.01	0.06	0.02
2-Methyl-1-butene (563-46-2)	52	8	0.02	0.03	0.01
2-Methyl-2-butene (513-35-9)	52	11	0.02	0.04	0.01
2-Methylheptane (592-27-8)	52	51	0.01	0.08	0.03
2-Methylhexane (591-76-4)	52	23	0.07	0.23	0.06
2-Methylpentane (107-83-5)	52	52	0.12	1.02	0.36
3-Methylheptane (589-81-1)	52	50	0.01	0.06	0.02
3-Methylhexane (589-34-4)	52	40	0.03	0.42	0.08
3-Methylpentane (96-14-0)	52	52	0.06	0.44	0.15
4-Methyl-1-pentene (691-37-2)	52	1	0.16	0.16	0.01
Acetylene (74-86-2)	52	51	0.10	0.82	0.32
a-Pinene (80-56-8)	52	17	0.01	0.10	0.02
Benzene (71-43-2)	52	51	0.06	0.33	0.14
cis-2-Butene (590-18-1)	52	18	0.01	0.05	0.01
cis-2-Hexene (7688-21-3)	52	1	0.01	0.01	0.01
cis-2-Pentene (627-20-3)	52	1	0.01	0.01	0.00
Cyclohexane (110-82-7)	52	52	0.06	0.62	0.21
Cyclopentane (287-92-3)	52	52	0.02	0.14	0.05
Cyclopentene (142-29-0)	52	10	0.03	0.09	0.05
Ethane (74-84-0)	52	52	5.30	76.50	23.69
Ethylbenzene (100-41-4)	52	24	0.01	0.03	0.01
Ethylene (74-85-1)	52	52	0.29	1.70	0.83
Isobutane (75-28-5)	52	52	0.42	7.68	2.44
Isopentane (78-78-4)	52	3	0.91	3.74	0.15
Isoprene (78-79-5)	52	33	0.01	0.45	0.06

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Bell Ranch (BRCO)
1/3/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
m-Diethylbenzene (141-93-5)	52	2	0.02	0.05	0.01
Methylcyclohexane (108-87-2)	52	49	0.14	0.86	0.30
Methylcyclopentane (96-37-7)	52	51	0.07	0.49	0.19
m-Ethyltoluene (620-14-4)	52	35	0.00	0.08	0.02
m-Xylene/p-Xylene (108-38-3/106-42-3)	52	52	0.02	0.12	0.05
n-Butane (106-97-8)	52	52	0.96	8.45	2.70
n-Decane (124-18-5)	52	34	0.01	0.13	0.02
n-Dodecane (112-40-3)	52	2	0.01	0.01	0.00
n-Heptane (142-82-5)	52	52	0.06	0.46	0.15
n-Hexane (110-54-3)	52	52	0.11	1.10	0.44
n-Nonane (111-84-2)	52	52	0.01	0.06	0.02
n-Octane (111-65-9)	52	49	0.04	0.21	0.07
n-Pentane (109-66-0)	52	52	0.25	5.20	1.14
n-Propylbenzene (103-65-1)	52	10	0.01	0.03	0.01
n-Tridecane (629-50-5)	52	1	0.02	0.02	0.00
n-Undecane (1120-21-4)	52	26	0.01	0.02	0.01
o-Ethyltoluene (611-14-3)	52	5	0.01	0.03	0.00
o-Xylene (95-47-6)	52	41	0.01	0.04	0.02
p-Diethylbenzene (105-05-5)	52	1	0.09	0.09	0.00
p-Ethyltoluene (622-96-8)	52	9	0.01	0.02	0.01
Propane (74-98-6)	52	52	2.07	31.63	9.87
Propylene (115-07-1)	52	52	0.06	0.46	0.22
Styrene (100-42-5)	52	3	0.05	0.52	0.01
Toluene (108-88-3)	52	52	0.09	0.66	0.21
trans-2-Butene (624-64-6)	52	22	0.01	0.21	0.03
trans-2-Pentene (646-04-8)	52	25	0.01	0.35	0.03

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
 Battlement Mesa (BMCO)
 1/21/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
1,2,3-Trimethylbenzene (526-73-8)	40	13	0.01	0.05	0.01
1,2,4-Trimethylbenzene (95-63-6)	40	35	0.01	0.11	0.04
1,3,5-Trimethylbenzene (108-67-8)	40	16	0.01	0.03	0.01
1,3-Butadiene (106-99-0)	40	2	0.01	0.01	0.01
1-Dodecene (112-41-4)	40	2	0.04	0.06	0.03
1-Hexene (592-41-6)	40	25	0.01	0.05	0.02
1-Nonene (124-11-8)	40	29	0.01	0.09	0.02
1-Octene (111-66-0)	40	37	0.01	0.11	0.03
1-Pentene (109-67-1)	40	37	0.01	0.08	0.03
1-Undecene (821-95-4)	40	6	0.01	0.02	0.03
2,2,3-Trimethylpentane (564-02-3)	40	13	0.01	0.04	0.01
2,2-Dimethylbutane (75-83-2)	40	37	0.01	0.05	0.03
2,3,4-Trimethylpentane (565-75-3)	40	11	0.01	0.03	0.01
2,3-Dimethylbutane (79-29-8)	40	40	0.02	0.08	0.06
2,3-Dimethylpentane (565-59-3)	40	40	0.01	0.06	0.04
2,4-Dimethylpentane (108-08-7)	40	39	0.01	0.09	0.03
2-Methyl-1-butene (563-46-2)	40	8	0.01	0.05	0.01
2-Methyl-2-butene (513-35-9)	40	15	0.02	0.05	0.02
2-Methylheptane (592-27-8)	40	40	0.02	0.09	0.05
2-Methylhexane (591-76-4)	40	20	0.04	0.25	0.07
2-Methylpentane (107-83-5)	40	39	0.11	0.63	0.33
3-Methylheptane (589-81-1)	40	40	0.01	0.08	0.04
3-Methylhexane (589-34-4)	40	34	0.03	0.21	0.09
3-Methylpentane (96-14-0)	40	40	0.05	0.21	0.14
Acetylene (74-86-2)	40	40	0.12	0.58	0.23
a-Pinene (80-56-8)	40	8	0.01	0.07	0.01
Benzene (71-43-2)	40	39	0.06	0.24	0.15
cis-2-Butene (590-18-1)	40	10	0.00	0.05	0.01
cis-2-Hexene (7688-21-3)	40	3	0.01	0.02	0.01
cis-2-Pentene (627-20-3)	40	3	0.01	0.10	0.01
Cyclohexane (110-82-7)	40	40	0.08	0.39	0.23
Cyclopentane (287-92-3)	40	40	0.02	0.07	0.04
Cyclopentene (142-29-0)	40	8	0.04	0.07	0.05
Ethane (74-84-0)	40	40	6.00	33.90	16.03
Ethylbenzene (100-41-4)	40	34	0.01	0.04	0.02
Ethylene (74-85-1)	40	40	0.40	1.60	0.89
Isobutane (75-28-5)	40	40	0.40	2.52	1.18
Isopentane (78-78-4)	40	1	0.70	0.70	0.02
Isoprene (78-79-5)	40	29	0.01	0.19	0.05
Isopropylbenzene (98-82-8)	40	5	0.01	0.02	0.01
m-Diethylbenzene (141-93-5)	40	7	0.01	0.13	0.02
Methylcyclohexane (108-87-2)	40	40	0.15	0.95	0.44

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
 Battlement Mesa (BMCO)
 1/21/2019-12/29/2019 (every sixth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
Methylcyclopentane (96-37-7)	40	40	0.07	0.31	0.20
m-Ethyltoluene (620-14-4)	40	27	0.01	0.08	0.02
m-Xylene/p-Xylene (108-38-3/106-42-3)	40	40	0.03	0.20	0.10
n-Butane (106-97-8)	40	40	0.50	2.80	1.33
n-Decane (124-18-5)	40	37	0.01	0.09	0.03
n-Dodecane (112-40-3)	40	5	0.01	0.02	0.01
n-Heptane (142-82-5)	40	40	0.07	0.37	0.18
n-Hexane (110-54-3)	40	40	0.11	0.74	0.37
n-Nonane (111-84-2)	40	40	0.01	0.10	0.04
n-Octane (111-65-9)	40	36	0.05	0.36	0.14
n-Pentane (109-66-0)	40	40	0.21	2.76	0.60
n-Propylbenzene (103-65-1)	40	10	0.01	0.05	0.01
n-Undecane (1120-21-4)	40	29	0.01	0.04	0.01
o-Ethyltoluene (611-14-3)	40	5	0.01	0.09	0.00
o-Xylene (95-47-6)	40	38	0.01	0.05	0.02
p-Diethylbenzene (105-05-5)	40	1	0.06	0.06	0.00
p-Ethyltoluene (622-96-8)	40	6	0.01	0.02	0.01
Propane (74-98-6)	40	40	1.70	10.73	4.94
Propylene (115-07-1)	40	40	0.08	0.66	0.25
Toluene (108-88-3)	40	40	0.12	0.59	0.32
trans-2-Butene (624-64-6)	40	12	0.04	0.13	0.03
trans-2-Pentene (646-04-8)	40	24	0.01	0.05	0.01

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
Carbondale (RFCO)
1/3/2019-12/23/2019 (every twelfth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
1,2,3-Trimethylbenzene (526-73-8)	26	8	0.01	0.05	0.01
1,2,4-Trimethylbenzene (95-63-6)	26	24	0.01	0.46	0.05
1,3,5-Trimethylbenzene (108-67-8)	26	7	0.01	0.03	0.01
1,3-Butadiene (106-99-0)	26	6	0.01	0.03	0.01
1-Dodecene (112-41-4)	26	3	0.02	0.16	0.03
1-Heptene (592-76-7)	26	15	0.01	0.05	0.02
1-Hexene (592-41-6)	26	14	0.01	0.05	0.01
1-Nonene (124-11-8)	26	21	0.01	0.03	0.02
1-Octene (111-66-0)	26	25	0.01	0.06	0.02
1-Pentene (109-67-1)	26	25	0.01	0.09	0.03
1-Undecene (821-95-4)	26	5	0.01	0.01	0.03
2,2,3-Trimethylpentane (564-02-3)	26	4	0.01	0.02	0.01
2,2,4-Trimethylpentane (540-84-1)	26	15	0.02	0.05	0.02
2,2-Dimethylbutane (75-83-2)	26	13	0.01	0.04	0.01
2,3,4-Trimethylpentane (565-75-3)	26	14	0.01	0.03	0.01
2,3-Dimethylbutane (79-29-8)	26	26	0.01	0.07	0.02
2,3-Dimethylpentane (565-59-3)	26	25	0.01	0.04	0.02
2,4-Dimethylpentane (108-08-7)	26	17	0.01	0.02	0.01
2-Methyl-1-butene (563-46-2)	26	9	0.02	0.05	0.02
2-Methyl-2-butene (513-35-9)	26	8	0.03	0.06	0.02
2-Methylheptane (592-27-8)	26	15	0.00	0.03	0.01
2-Methylhexane (591-76-4)	26	9	0.04	0.13	0.03
2-Methylpentane (107-83-5)	26	25	0.06	0.43	0.15
3-Methylheptane (589-81-1)	26	11	0.01	0.02	0.01
3-Methylhexane (589-34-4)	26	9	0.03	0.30	0.03
3-Methylpentane (96-14-0)	26	25	0.02	0.15	0.05
Acetylene (74-86-2)	26	25	0.11	1.44	0.56
a-Pinene (80-56-8)	26	21	0.01	0.40	0.08
Benzene (71-43-2)	26	26	0.06	0.21	0.11
b-Pinene (127-91-3)	26	1	0.06	0.06	0.01
cis-2-Butene (590-18-1)	26	18	0.01	0.03	0.01
cis-2-Hexene (7688-21-3)	26	1	0.02	0.02	0.01
cis-2-Pentene (627-20-3)	26	5	0.00	0.01	0.00
Cyclohexane (110-82-7)	26	26	0.02	0.23	0.05
Cyclopentane (287-92-3)	26	23	0.01	0.05	0.02
Cyclopentene (142-29-0)	26	5	0.04	0.06	0.05
Ethane (74-84-0)	26	26	1.75	25.20	4.75
Ethylbenzene (100-41-4)	26	25	0.01	0.05	0.02
Ethylene (74-85-1)	26	26	0.27	2.06	0.99
Isobutane (75-28-5)	26	26	0.17	2.40	0.49
Isopentane (78-78-4)	26	2	0.26	0.27	0.02
Isoprene (78-79-5)	26	18	0.01	1.03	0.15

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

Garfield County SNMOC Monitoring
 Carbondale (RFCO)
 1/3/2019-12/23/2019 (every twelfth day)

Detected Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
Isopropylbenzene (98-82-8)	26	3	0.01	0.02	0.01
m-Diethylbenzene (141-93-5)	26	4	0.03	0.10	0.02
Methylcyclohexane (108-87-2)	26	13	0.02	0.31	0.03
Methylcyclopentane (96-37-7)	26	24	0.03	0.18	0.06
m-Ethyltoluene (620-14-4)	26	24	0.02	0.16	0.04
m-Xylene/p-Xylene (108-38-3/106-42-3)	26	26	0.02	0.10	0.06
n-Butane (106-97-8)	26	26	0.34	2.65	0.95
n-Decane (124-18-5)	26	19	0.01	0.06	0.02
n-Dodecane (112-40-3)	26	1	0.01	0.01	0.00
n-Heptane (142-82-5)	26	26	0.02	0.14	0.04
n-Hexane (110-54-3)	26	26	0.06	0.37	0.14
n-Nonane (111-84-2)	26	23	0.01	0.04	0.01
n-Octane (111-65-9)	26	26	0.01	0.07	0.03
n-Pentane (109-66-0)	26	26	0.13	5.42	0.54
n-Propylbenzene (103-65-1)	26	7	0.01	0.06	0.01
n-Undecane (1120-21-4)	26	9	0.01	0.02	0.01
o-Ethyltoluene (611-14-3)	26	9	0.01	0.03	0.01
o-Xylene (95-47-6)	26	25	0.01	0.05	0.02
p-Diethylbenzene (105-05-5)	26	1	0.03	0.03	0.00
p-Ethyltoluene (622-96-8)	26	8	0.01	0.06	0.01
Propane (74-98-6)	26	26	1.97	10.07	4.36
Propylene (115-07-1)	26	26	0.08	0.74	0.24
Propyne (74-99-7)	26	1	0.02	0.02	0.00
Styrene (100-42-5)	26	3	0.07	0.43	0.03
Toluene (108-88-3)	26	24	0.11	1.69	0.43
trans-2-Butene (624-64-6)	26	6	0.01	0.08	0.01
trans-2-Hexene (4050-45-7)	26	1	0.10	0.10	0.01
trans-2-Pentene (646-04-8)	26	23	0.01	0.03	0.02

*Samples reported as non-detects (ND) were included in averages as 1/2 minimum detection limits.

APPENDIX C

Carbonyl Concentrations

Garfield County Carbonyl Monitoring
Parachute (PACO)
1/3/2019-11/23/2019 (every twelfth day)

Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
2,5-Dimethylbenzaldehyde (5779-94-2)	7	0	ND	ND	0.00
2-Butanone (78-93-3)	23	17	0.03	0.11	0.06
Acetaldehyde (75-07-0)	23	23	0.02	0.85	0.37
Acetone (67-64-1)	23	23	0.14	2.07	1.17
Benzaldehyde (100-52-7)	23	19	0.01	0.10	0.02
Butyraldehyde (123-72-8)	23	18	0.01	0.04	0.02
Crotonaldehyde (123-73-9)	23	18	0.01	0.14	0.03
Formaldehyde (50-00-0)	23	23	0.03	2.37	1.18
Hexaldehyde (66-25-1)	23	15	0.01	0.22	0.04
Isovaleraldehyde (590-86-3)	7	0	ND	ND	0.01
Propionaldehyde (123-38-6)	23	18	0.01	0.07	0.03
Tolualdehydes (NA)	7	2	0.03	0.03	0.01
Valeraldehyde (110-62-3)	23	10	0.01	0.02	0.01

*Samples reported as non-detects (ND) are included in averages as 1/2 minimum detection limits.

Garfield County Carbonyl Monitoring
Rifle Library (RECO)
1/3/2019-12/29/2019 (every twelfth day)

Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
2,5-Dimethylbenzaldehyde (5779-94-2)	8	0	ND	ND	0.00
2-Butanone (78-93-3)	27	21	0.01	0.69	0.09
Acetaldehyde (75-07-0)	27	26	0.01	0.96	0.44
Acetone (67-64-1)	27	27	0.07	2.11	1.03
Benzaldehyde (100-52-7)	27	17	0.01	0.10	0.02
Butyraldehyde (123-72-8)	27	17	0.01	0.05	0.02
Crotonaldehyde (123-73-9)	27	18	0.01	0.16	0.03
Formaldehyde (50-00-0)	27	27	0.01	2.16	0.95
Hexaldehyde (66-25-1)	27	21	0.01	0.26	0.05
Isovaleraldehyde (590-86-3)	8	0	ND	ND	0.00
Propionaldehyde (123-38-6)	27	23	0.01	0.09	0.03
Tolualdehydes (NA)	8	2	0.02	0.04	0.01
Valeraldehyde (110-62-3)	27	10	0.01	0.02	0.01

*Samples reported as non-detects (ND) are included in averages as 1/2 minimum detection limits.

Garfield County Carbonyl Monitoring
 Bell Ranch (BRCO)
 1/3/2019-10/6/2019 (every twelfth day)

Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
2,5-Dimethylbenzaldehyde (5779-94-2)	7	0	ND	ND	0.00
2-Butanone (78-93-3)	19	13	0.01	0.13	0.05
Acetaldehyde (75-07-0)	19	19	0.01	0.44	0.19
Acetone (67-64-1)	19	19	0.04	1.27	0.58
Benzaldehyde (100-52-7)	19	12	0.01	0.04	0.01
Butyraldehyde (123-72-8)	19	8	0.01	0.04	0.01
Crotonaldehyde (123-73-9)	19	8	0.01	0.12	0.02
Formaldehyde (50-00-0)	19	19	0.02	1.14	0.45
Hexaldehyde (66-25-1)	19	11	0.02	0.14	0.02
Isovaleraldehyde (590-86-3)	7	0	ND	ND	0.00
Propionaldehyde (123-38-6)	19	13	0.01	0.06	0.02
Tolualdehydes (NA)	7	1	0.04	0.04	0.01
Valeraldehyde (110-62-3)	19	3	0.01	0.02	0.00

*Samples reported as non-detects (ND) are included in averages as 1/2 minimum detection limits.

Garfield County Carbonyl Monitoring
 Battlement Mesa (BMCO)
 1/3/2019-12/17/2019 (every twelfth day)

Compound (CAS Number)	Sample Count		Concentration (ppbV)		
	# Samples	# Detects	Minimum	Maximum	Average*
2,5-Dimethylbenzaldehyde (5779-94-2)	7	0	ND	ND	0.00
2-Butanone (78-93-3)	28	21	0.02	0.12	0.07
Acetaldehyde (75-07-0)	28	28	0.02	0.80	0.35
Acetone (67-64-1)	28	28	0.09	2.41	1.09
Benzaldehyde (100-52-7)	28	23	0.01	0.07	0.02
Butyraldehyde (123-72-8)	28	21	0.01	0.04	0.02
Crotonaldehyde (123-73-9)	28	21	0.01	0.15	0.03
Formaldehyde (50-00-0)	28	28	0.02	1.93	0.96
Hexaldehyde (66-25-1)	28	22	0.01	0.15	0.03
Isovaleraldehyde (590-86-3)	7	0	ND	ND	0.00
Propionaldehyde (123-38-6)	28	26	0.01	0.09	0.03
Tolualdehydes (NA)	7	3	0.02	0.03	0.01
Valeraldehyde (110-62-3)	28	12	0.01	0.03	0.01

*Samples reported as non-detects (ND) are included in averages as 1/2 minimum detection limits.